

A CLASS OF SPACE-TIMES OF NON-RIGIDLY ROTATING DUST

ŁUKASZ BRATEK

*Institute of Nuclear Physics PAN, Radzikowskiego 152, 31-342
Kraków, Poland. e-mail: lukasz.brateg@ifj.edu.pl*

JOANNA JAŁOCHA

*Jagellonian University, Institute of Physics, Reymonta 4,
30-059 Kraków, Poland. e-mail:*

MAREK KUTSCHERA

*Institute of Nuclear Physics PAN, Radzikowskiego 152,
31-342 Kraków, Poland, & Jagellonian University, Institute
of Physics, Reymonta 4, 30-059 Kraków, Poland e-mail:
marek.kutschera@ifj.edu.pl*

Abstract. We find a class of exact solutions of differentially rotating dust in the framework of General Relativity. There exist asymptotically flat space-times of the flow with positive mass function that for radii sufficiently large is monotonic and tends to zero at infinity. Some of the space-times may have non-vanishing total angular momentum. The flow is essentially different from another exactly solvable flow described by van Stockum line element.

1. INTRODUCTION

The paper is devoted to investigation of solutions of self-gravitating cylindrically symmetric dust flow on circular orbits along field lines of locally non-rotating observers in the framework of General Relativity. The flow is non-rigid (differential) and non-expanding. It turns out the resulting space-times are globally regular, apart from the points where singularities may be located, and that the congruence of locally non-rotating observers used to define energy-momentum exist globally, as well. Consequently, the problem of the flow is well posed. We shall call such a flow the $\mathcal{K}$ **flow** for brevity. The main drawback of the flow is that the proper energy density is negative definite and that the flow is purely relativistic – has no newtonian limit.

The result of our paper is the construction of the multipole sequence of asymptotically flat, stationary, and cylindrically symmetric external solutions to which a broad class of other asymptotically flat solutions can be decomposed. We construct also the corresponding multipole sequence of internal space-times. The multipole representation is enumerable by a discrete parameter. There exist also a continuum spectrum of basic solutions of which we do not examine here as they are nonanalytic on the axis of rotation. The stationary part of multipolar solutions that are asymptotically flat we call external multipoles, and the stationary part of multipolar solutions that are not asymptotically flat we call internal multipoles.

We showed that external multipoles, and any asymptotically flat space-times of $\mathcal{K}$ flow, are massless no matter what is pasted smoothly as an alternative internal completion of the space-times. Despite the fact, some of the space-times can have non-vanishing total angular momentum.

As in standard gravity a broad class of solutions can be decomposed to the multipoles. Even though the resulting space-times are solutions of Einstein equations of a differentially rotating dust, thus physically more interesting than van Stockum–Bonnor flows [1], [3], [4], it seems that they are of no physical concern as local energy density is not positive definite. Anyway, we decided to publish the solutions as they present another class of exactly solvable dust flows in general relativity. Similarly, as the exactly solvable van Stockum–Bonnor class of asymptotically flat flows along time Killing vector field, the asymptotically flat external multipoles of $\mathcal{K}$ flows, which must have vanishing total mass irrespectively of internal deformations¹, have no newtonian limit and are useless in astrophysical situations. As far as exotic forms of matter concern, the internal solutions of both the models can be still considered as physically interesting and worth to be examined if completed smoothly by, say, asymptotically flat external (Kerr) vacuum solutions.

It turns out that the structure function $K = \xi\eta$ of the flow is the same as for the van-Stockum Bonnor flow [4], nonetheless, both the flows and geometry of the resulting space-times are qualitatively different. An asymptotically flat van Stockum–Bonnor flow is rigid, non-expanding, and has positive energy density proportional to the square of the vorticity scalar which does not vanish. An asymptotically flat $\mathcal{K}$ flow is differential, non-expanding, locally non-rotating, and has negative definite energy density proportional to the square of the non-vanishing shear tensor. Dust of van Stockum–Bonnor does not rotate with respect to asymptotic observers, it has a point-dependent physical velocity as measured with respect to the locally dragged inertial frames and which is proportional to K , whereas dust of a $\mathcal{K}$ flow rotates differentially with respect to asymptotic observers with angular velocity proportional to K , and has vanishing physical velocity. The peculiarities are possible due to purely relativistic effects and show the flows have no newtonian limits.

As an example we constructed a z -symmetric, cylindrically symmetric, and stationary for radii sufficiently large, asymptotically flat space-time which is globally smooth (with the exception of point-like singularities of curvature residing on the axis of rotation), and which can be decomposed globally to the external multipoles. The solution has nonzero total angular momentum and zero total mass.

2. SETUP

We assume the global existence of the cylindrical symmetry space-like Killing vector η with closed field lines, together with the global existence of time translation Killing vector ξ of which field lines are opened, time-like for radii sufficiently large, and asymptotically normalizable to unity. Having in mind stationary and asymptotically flat space-times with cylindrical symmetry, we shall proceed as in the standard theory of rotating stars in the relativistic astrophysics [2]. The assumption of asymptotical flatness allows for the unique determination of Killing vectors ξ and η by the defining properties. In addition one assumes that asymptotically $\xi\eta \rightarrow 0$. Field lines of the Killing vectors, which clearly are frame independent

¹one can imagine such a deformation as a replacement of the interior by a region of arbitrary space-time such the resulting space-time be still smooth

objects, may be viewed as two of four coordinate lines in some particular coordinates in which the time coordinate t runs along open lines of ξ and the cyclic coordinate ϕ along closed lines of η , that is, $\xi \equiv \partial_t$ and $\eta \equiv \partial_\phi$ by definition. The other two, denoted by ρ and z , are arbitrary internal coordinates in a two dimensional subspace orthogonal to ξ and η . In this coordinates the most general line element of a cylindrically symmetric and stationary space-time is fully determined by four structure functions $\nu(\rho, z)$, $\psi(\rho, z)$, $\tilde{\omega}(\rho, z)$ and $\mu(\rho, z)$ and reads [2]

$$(2.1) \quad ds^2 = e^{2\lambda} dt^2 - e^{2\psi} (d\phi - \tilde{\omega} dt)^2 - e^{2\mu} (d\rho^2 + dz^2).$$

Bardeen coordinates are distinguished by the property that in these coordinates Killing vectors ξ and η attain particularly simple form

$$\xi^\mu = \delta_t^\mu, \quad \eta^\mu = \delta_\phi^\mu.$$

In asymptotically flat space-time one can introduce cylindrical coordinate system in which the line element at infinity reduces to

$$ds^2 \rightarrow dt^2 - d\rho^2 - \rho^2 d\phi^2 - dz^2$$

Axis of the coordinates frame are attached to 'fixed stars', otherwise the axis would rotate. This in turn would be in contradiction with the assumption that ∂_t can be asymptotically normalized to unity. One assumes, therefore, that asymptotically condition $e^{2\lambda} - \tilde{\omega}^2 e^{2\psi} > 0$ should hold. The latter is not the case e.g. in a rotating frame of reference that rotates with constant angular velocity. Put differently, Bardeen coordinates are asymptotically inertial (anyway, this metric can be treated more formally and be used to describe solutions that are not asymptotically flat).

As it turns out later, $\mathcal{K}$ flow can be defined globally, that is, locally non-rotating observers exist for all points (apart from singularities), but, unlike for van-Stockum Bonnor flow, space-time of $\mathcal{K}$ flow is not globally stationary. The $\mathcal{K}$ flow is differential (the shear tensor does not vanish). The angular velocity of asymptotically flat $\mathcal{K}$ flow is, by construction, identical to the angular velocity of dragging of inertial frames $\tilde{\omega}$. As so, dust rotates on circular orbits with respect to asymptotical stationary observers even though it locally does not rotate (the vorticity vector of the flow vanishes identically) and its physical velocity, measured with respect to dragged local inertial frames, is also identically zero.

2.1. The line element of $\mathcal{K}$ flow. We derive the line element of the $\mathcal{K}$ flow step by step from the general form of the line element given by 2.1. We specify the structure functions uniquely by the requirement that dust space-time trajectories are identical with field lines of four-velocity field of locally non-rotating observers and that Einstein equations are satisfied.

2.1.1. A note on locally non-rotating observers. It should be clear that irrespectively of any particular reference frame, a generic cylindrically symmetric and stationary flow is fully determined by a four-velocity field

$$U(\Omega) = Z(\Omega) (\xi + \Omega \eta),$$

provided

$$Z(\Omega)^{-2} \equiv \xi\xi + 2\Omega\xi\eta + \Omega^2\eta\eta > 0, \quad \mathcal{L}_\xi\Omega = 0 = \mathcal{L}_\eta\Omega.$$

In an asymptotically flat space-time Ω has the interpretation of the angular velocity measured with respect to 'fixed stars'. In particular, the condition $\mathbf{U}(\Omega)\boldsymbol{\eta} = 0$ gives

$$\tilde{\Omega} = -\frac{\boldsymbol{\xi}\boldsymbol{\eta}}{\boldsymbol{\eta}\boldsymbol{\eta}} \equiv \tilde{\omega}.$$

The vorticity tensor² of velocity field $\mathbf{n} \equiv \mathbf{U}(\tilde{\omega})$ vanishes identically, which can be checked by a direct calculation in Bardeen coordinates 2.1. Therefore, the observers are called locally non-rotating, although they move differentially on circular orbits with respect to 'fixed stars' with angular velocity $\tilde{\omega}$ of dragging of inertial frames. Indeed, congruence of locally non-rotating observers distorts without changing proper volume³

$$\Omega^2(\mathbf{n}) \equiv 0, \quad \Theta(\mathbf{n}) \equiv 0, \quad \sigma^2(\mathbf{n}) = -\frac{1}{4} \frac{(\boldsymbol{\eta}\boldsymbol{\eta})^2 (\nabla\tilde{\omega})^2}{\begin{vmatrix} \boldsymbol{\xi}\boldsymbol{\eta} & \boldsymbol{\xi}\boldsymbol{\xi} \\ \boldsymbol{\eta}\boldsymbol{\eta} & \boldsymbol{\xi}\boldsymbol{\eta} \end{vmatrix}}, \quad \tilde{\omega} = -\frac{\boldsymbol{\xi}\boldsymbol{\eta}}{\boldsymbol{\eta}\boldsymbol{\eta}}.$$

Local inertial frames tangent to $\mathbf{n}$ play the role of local standards of rest with respect to which 'physical' velocity is measured. Therefore, the physical velocity of a particle moving with four-velocity $\mathbf{U}(\Omega)$ is the relative velocity of two inertial observers momentarily co-moving with $\mathbf{U}(\Omega)$ and $\mathbf{n}$, respectively. After Special Relativity we define the relative velocity as the hyperbolic tangent of the hyperbolic angle χ between four-velocities $\mathbf{U}(\Omega)$ and $\mathbf{n}$. In Bardeen coordinates it reads

$$(2.2) \quad \tanh \chi = e^{\psi-\lambda}(\Omega - \tilde{\omega}).$$

The result agrees with that of [2] where more 'physical' derivation was presented.

2.1.2. *Specifying the line element.* The energy-momentum tensor of dust matter moving along field lines of vector field $\mathbf{n}$ reads $\mathbf{T} = \mathcal{D}\mathbf{U} \otimes \mathbf{U}$ where

$$\mathbf{U}(\tilde{\omega}) = Z(\tilde{\omega})(\boldsymbol{\xi} + \tilde{\omega}\boldsymbol{\eta}) \quad \Rightarrow \quad U^\mu = e^{-\lambda}[1, 0, \tilde{\omega}, 0].$$

From the previous construction it follows that dust in asymptotically flat $\mathcal{K}$ flow moves differentially on circular orbits with angular velocity of dragging of inertial frames $\tilde{\omega}$ as seen by an asymptotic stationary observer. This flow has vanishing vorticity vector, that is, it does not rotate locally, and has no physical velocity – it is at rest with respect to congruence of local standards of rest dragged with angular velocity $\tilde{\omega}$. It follows also that angular momentum per mass element of the flow $-U^\mu \eta_\mu$ is identically zero as $\mathbf{U} \equiv \mathbf{n}$ and $\boldsymbol{\eta}$ are orthogonal. Nevertheless, as we shall show later, total angular momentum does not vanish for particular kinds of asymptotically flat space-times of $\mathcal{K}$ flow which must contain singularities that contribute total angular momentum of the space-times.

Einstein's equations $\mathbf{G} = \kappa\mathbf{T}$ and the contracted Bianchi identities $\nabla^\mu G_{\mu\nu} = 0$ yield local conservation law $\nabla^\mu T_{\mu\nu} = 0$ that for dust gives continuous flow along

²see footnote 3

³Dilation tensor $\Theta(\mathbf{u})$ and (traceless) shear tensor $\sigma(\mathbf{u})$ are defined for a velocity field $\mathbf{u}$ as $\Theta_{\mu\nu} = \nabla_\alpha u^\alpha h_{\mu\nu}$ and $\sigma_{\mu\nu} = \nabla_{(\alpha} u_{\beta)} h^\alpha_\mu h^\beta_\nu - \frac{1}{3} \nabla_\alpha u^\alpha h_{\mu\nu}$, where $h^\mu_\nu = \delta^\mu_\nu - u^\mu u_\nu$ is projector $\mathbf{h}(\mathbf{u})$ onto subspace orthogonal to $\mathbf{u}$. Vorticity tensor $\Omega(\mathbf{u})$ for $\mathbf{u}$ is defined as $\omega_{\mu\nu} = \nabla_{[\alpha} u_{\beta]} h^\alpha_\mu h^\beta_\nu$ and it yields a derivative quantity that characterizes vertex – the vorticity vector $\omega^\mu = \frac{1}{2} \frac{\epsilon^{\mu\alpha\beta\gamma}}{\sqrt{-g}} u_\alpha \omega_{\beta\gamma}$. One defines also square of dilation scalar $\sigma^2(\mathbf{u}) = \frac{1}{2} \sigma^{\mu\nu} \sigma_{\mu\nu} \geq 0$ and square of vorticity scalar $\Omega^2(\mathbf{u}) = \frac{1}{2} \omega^{\mu\nu} \omega_{\mu\nu} \geq 0$, then $\Omega^2(\mathbf{u}) = -\omega^\mu \omega_\mu$.

geodesic paths

$$\nabla_\mu (\mathcal{D}U^\mu) = 0 \quad \text{and} \quad U^\nu \nabla_\nu U_\mu = 0.$$

The continuity equation is satisfied identically for $\mathcal{K}$ flow on the power of Killing equations and of symmetries of energy density $\xi\mathcal{D} = 0$ and $\eta\mathcal{D} = 0$. In Bardeen coordinates $U^\nu \nabla_\nu U_\mu = \{0, -\partial_\rho \lambda, 0, -\partial_z \lambda\}$ which implies that λ must be a constant. Another constraint on structure functions follows from the linear combination of Einstein's equations

$$0 \equiv \xi^{[\mu} \eta^{\alpha]} \xi_{[\alpha} \eta_{\nu]} \left(R_{\mu}{}^{\nu} - \kappa \left(T_{\mu}{}^{\nu} - \frac{1}{2} T \delta_{\mu}{}^{\nu} \right) \right) = \frac{1}{4} e^{\lambda+\psi-2\mu} (\partial_\rho^2 + \partial_z^2) e^{\lambda+\psi}.$$

The left side is identically zero on the power of Einstein equations, while the expression to the right holds for pressureless perfect fluid in Bardeen coordinates. We have not yet specified the arbitrary coordinates ρ and z . Asymptotical flatness in cylindrical coordinates requires that $e^{2\lambda} \rightarrow 1$ and $e^{2\psi} \rightarrow \rho^2$, thus $e^{\lambda+\psi} \rightarrow \rho$. The simplest way to concord the facts is to choose as a solution of the above equation

$$(2.3) \quad \lambda = 0, \quad e^\psi = \rho.$$

We have thus shown that $\mathcal{K}$ flow requires at most two structure functions $\tilde{\omega}$ and μ .

Killing vector η is always space-like $\eta\eta \equiv -\rho^2$ thus the space-time of $\mathcal{K}$ flow is globally cylindrically symmetric. As $\xi\xi = 1 - \tilde{\omega}^2 \rho^2$ it is also stationary in regions where $\tilde{\omega}^2 \rho^2 < 1$. Note that $\det[g] = -\rho^2 e^{4\mu}$ does not depend explicitly on $\tilde{\omega}$ thus the boundary surface $\tilde{\omega}^2 \rho^2 = 1$, which demarcates stationary and non-stationary regions of a space-time, is nonsingular, and in a sense, is alike to Schwarzschild horizon.

We stress the construction of the energy-momentum tensor of $\mathcal{K}$ flow makes sense globally as $U^\mu U_\mu > 0$, or $\xi\xi + 2\tilde{\omega}\xi\eta + \tilde{\omega}^2\eta\eta \equiv 1 > 0$ for $\mathcal{K}$ flow. $U(\tilde{\omega})$ is time-like even though in some regions the resulting space-time may be non-stationary, and so, $U(\tilde{\omega})$ be a combination of two space-like vectors. It follows that in space-times of $\mathcal{K}$ flow the locally non-rotating observers exist globally.

Similarly as in [4] we define $K = \xi\eta$ then in Bardeen coordinates $K = \rho^2 \tilde{\omega}$, and (K, μ) can be used in place of $(\tilde{\omega}, \mu)$ as structure functions of $\mathcal{K}$ flow. We have thus reduced the line element 2.1 of $\mathcal{K}$ flow to

$$(2.4) \quad ds^2 = \left(1 - \frac{K^2(\rho, z)}{\rho^2} \right) dt^2 + 2K(\rho, z) dt d\phi - \rho^2 d\phi^2 - e^{2\mu(\rho, z)} (d\rho^2 + dz^2).$$

We have seen, that in space-times given by the line element, world-lines of locally non-rotating observers are geodesics.

At a given fixed point one can diagonalize the line element by a local linear transformation (local boost) of the base forms, say, $d\phi \rightarrow d\phi + \tilde{\omega} dt$. At the fixed point the line element reduces simply to $dt^2 - \rho^2 d\phi^2 - e^{2\mu} (d\rho^2 + dz^2)$. The new base overlaps with the inertial frame of an observer momentarily co-moving with the locally non-rotating observer at that point.

2.2. Equations for structure functions of $\mathcal{K}$ flow. We have shown that for $\mathcal{K}$ flow general metric 2.1 can be reduced to 2.4. In what follows we shall derive equations for K and μ . Let $E = G - \kappa T$ with $T = \mathcal{D}U \otimes U$, $U = n$. Einstein's equation $E = 0$ imply from

$$E^\rho{}_\rho = \frac{e^{-2\mu}}{\rho} \left(\rho^3 \frac{\tilde{\omega}_{,z}^2 - \tilde{\omega}_{,\rho}^2}{4} - \mu_{,\rho} \right), \quad E^\rho{}_z = -\frac{e^{-2\mu}}{\rho} \left(\rho^3 \frac{\tilde{\omega}_{,\rho} \tilde{\omega}_{,z}}{2} + \mu_{,z} \right)$$

($E^\rho_\rho = -E^z_z$, $E^\rho_z = E^z_\rho$) that

$$(2.5) \quad \mu_{,\rho} = \rho^3 \frac{\tilde{\omega}_{,z}^2 - \tilde{\omega}_{,\rho}^2}{4}, \quad \mu_{,z} = -\rho^3 \frac{\tilde{\omega}_{,\rho} \tilde{\omega}_{,z}}{2}.$$

For $\mathcal{C}^2$ solutions the Schwarz identity $\mu_{,\rho z} = \mu_{,z\rho}$ imposes on $\tilde{\omega}$ the linear elliptic constraint

$$(2.6) \quad \tilde{\omega}_{,\rho\rho} + 3\rho^{-1}\tilde{\omega}_{,\rho} + \tilde{\omega}_{,zz} = 0 \quad \Leftrightarrow \quad K_{,\rho\rho} - \frac{K_{,\rho}}{\rho} + K_{,zz} = 0.$$

As

$$E^t_\phi = \frac{e^{-2\mu}}{2} \left(K_{,\rho\rho} - \frac{K_{,\rho}}{\rho} + K_{,zz} \right)$$

equation $E^t_\phi = 0$ is satisfied identically. By calculating $\mu_{,\rho\rho}$ and $\mu_{,zz}$ from 2.5 and using 2.6, one may check the component E^t_t (then $E^\phi_t = K\rho^{-2}E^t_t$) reduces to $E^t_t = -e^{-2\mu}\rho^2(\tilde{\omega}_{,\rho}^2 + \tilde{\omega}_{,z}^2) - \kappa\mathcal{D} = 0$, and finally,

$$(2.7) \quad \kappa\mathcal{D} = -e^{-2\mu}\rho^2(\tilde{\omega}_{,\rho}^2 + \tilde{\omega}_{,z}^2).$$

The other components of $\mathbf{E}$ vanish identically by symmetry. As energy density is negative definite $\mathcal{K}$ flow is unphysical in the sense it cannot be made of ordinary matter. Once a solution of 2.6 is found, which is simple task due to linearity, 2.7 gives the respective energy density and 2.5 can be easily integrated. Contribution to the total mass, from regions where $\tilde{\omega} \in \mathcal{C}^1$, is negative and given for solutions by

$$(2.8) \quad M_{smooth} = -\frac{c^2}{4G} \int \rho^3(\tilde{\omega}_{,\rho}^2 + \tilde{\omega}_{,z}^2) d\rho dz$$

anyway, we shall show that asymptotically flat space-times of $\mathcal{K}$ flow must contain distributional sources of a nett positive mass $-M_{smooth}$ that cancels contributions (possibly infinite) from smooth components given by integrals as the above, as the asymptotic total mass, calculated as a surface integral at infinity, vanishes for asymptotically flat $\mathcal{K}$ flows.

Note, that elliptic equation 2.6 is the condition for critical points $\tilde{\omega}$ of energy functional 2.8 (when derived by integrating 2.7 with respect to the proper volume $e^{2\mu}\rho$ of the surface of constant time, then $M_{smooth} = \int \mathcal{D}e^{2\mu}\rho d\rho d\phi dz$) in any finite region of a hyper-surface of constant time bounded by a closed line in the plane (ρ, z) and such that $\tilde{\omega}$ is at least $\mathcal{C}^1$ (piecewise) in the region, namely

$$\delta M[\tilde{\omega}] = 0 \quad \Rightarrow \quad \tilde{\omega}_{,\rho\rho} + 3\frac{\tilde{\omega}_{,\rho}}{\rho} + \tilde{\omega}_{,zz} = 0.$$

For the solutions the maximum (minimum) principle applies.

2.3. Some solutions. To obtain basic solutions we transform 2.6 to spherical coordinates $\rho \rightarrow r \sin \theta$, $z \rightarrow r \cos \theta$. By substituting $K(r, \theta) = R(r)Y(\theta)$, the separation of variables gives $R(r) = r^{-n}$ or $R(r) = r^{n+1}$, and the hypergeometric equation for $Y(x)$, where $x = \cos \theta$. We assume here $n \in \mathbb{N}$ and take only solutions that are analytic at $x = \pm 1$, by which the multipole series is established. The general

formula for z -antisymmetric and z -symmetric external multipoles that are solutions of 2.6 and that give asymptotically flat space-times is

$$(2.9) \quad K(\rho, z) = \begin{cases} \frac{1}{(\rho^2 + z^2)^{m-1/2}} {}_2F_1\left(m - \frac{1}{2}, -m; \frac{1}{2}, \frac{z^2}{\rho^2 + z^2}\right), & m = 1, 2, 3, \dots \\ \frac{z}{(\rho^2 + z^2)^{m+1/2}} {}_2F_1\left(m + \frac{1}{2}, -m; \frac{3}{2}, \frac{z^2}{\rho^2 + z^2}\right), & m = 1, 2, 3, \dots \end{cases}$$

For $m = 0$ one obtains $K = \sqrt{\rho^2 + z^2}$ and the monopole⁴ $K = z/\sqrt{\rho^2 + z^2}$ which are non-flat asymptotically. The corresponding series of internal multipoles (singular at infinity) is

$$(2.10) \quad K(\rho, z) = \begin{cases} (\rho^2 + z^2)^m {}_2F_1\left(m - \frac{1}{2}, -m; \frac{1}{2}, \frac{z^2}{\rho^2 + z^2}\right), & m = 0, 1, 2, \dots \\ z(\rho^2 + z^2)^m {}_2F_1\left(m + \frac{1}{2}, -m; \frac{3}{2}, \frac{z^2}{\rho^2 + z^2}\right), & m = 0, 1, 2, \dots \end{cases}$$

2.3.1. *An example.* In the previous paper we found an example of asymptotically flat solution (note that equation for K function are identical but the models and line elements are essentially different)

$$(2.11) \quad K_a(\rho, z) = \frac{2\rho^2 z}{(z+a)\sqrt{\rho^2 + (z-a)^2} + (z-a)\sqrt{\rho^2 + (z+a)^2}} + \dots \\ \dots + \frac{\rho^2}{2a} \ln \left(\frac{z-a + \sqrt{\rho^2 + (z-a)^2}}{z+a + \sqrt{\rho^2 + (z+a)^2}} \right)$$

which can be expanded in the series of external z -symmetric multipoles

$$K_a(\rho, z) = \frac{2}{3} \frac{\rho^2}{r^3} a^2 - \frac{1}{5} \frac{\rho^2 (\rho^2 - 4z^2)}{r^7} a^4 + \frac{3}{28} \frac{\rho^2 (\rho^4 - 12\rho^2 z^2 + 8z^4)}{r^{11}} a^6 + \dots,$$

where $r = \sqrt{\rho^2 + z^2}$. Solution K_a has single extremum $K(0, 0) = a$ and, apart from the singular points $(0, a)$ and $(0, -a)$, is everywhere smooth. The two points are singular since, for example, $\partial_z K$ as a function of z , for $\rho = 0$ has a jump at $z = \pm a$, $a > 0$. The corresponding angular velocity $\tilde{\omega}_a = \rho^{-2} K_a$ has additional singularities, namely it is analytical everywhere with the exception of the closed segment $\rho = 0$, $z \in [-a, a]$. The resulting space-time is asymptotically flat

$$K_a(r, \theta) \sim \frac{2}{3} \frac{a^2}{r} \sin^2 \theta, \quad \tilde{\omega}_a(r, \theta) \sim \frac{2}{3} \frac{a^2}{r^3}, \quad \mu_a(r, \theta) \sim \frac{a^4 \sin^4 \theta}{4r^4}, \quad r \rightarrow \infty,$$

and

$$g_{tt} \sim 1 - \frac{4}{9} \frac{a^4}{r^4} \sin^2 \theta, \quad g_{t\phi} \sim \frac{2}{3} \frac{a^2}{r} \sin^2 \theta \left(1 + \frac{3}{20} \frac{a^2}{r^2} (3 + 5 \cos 2\theta) \right), \\ g_{\phi\phi} = -r^2 \sin^2 \theta, \quad g_{rr} = \frac{g_{\theta\theta}}{r^2} \sim -1 - \frac{a^4}{2r^4} \sin^4 \theta.$$

Comparison with asymptotical expansion of the Kerr metric allows to determine total mass of the space-time, which is $M = 0$, and total angular momentum, which is $J = a^2/3$. The result proves that the singular segment of solution $\tilde{\omega}_a$, located on the z -axis at $-a \leq z \leq a$, are sources of positive mass that cancel negative mass of the remaining regions ($\mathcal{D} < 0$). In general, total mass of asymptotically

⁴We assume the same nomenclature for the K as in [4] where Euclidean mass density $\rho^{-2} (K_{,\rho}^2 + K_{,z}^2)$ for the monopole of the van Stockum-Bonnor flow was spherically symmetric, even though, as one can check, the Euclidean mass density $-\rho^{+2} (\tilde{\omega}_{,\rho}^2 + \tilde{\omega}_{,z}^2)$ for the monopole $\tilde{\omega} = K/\rho^2$ of the $\mathcal{K}$ flow is not spherically symmetric.

flat space-time can be calculated as surface integral over sphere $\mathcal{S}^2$ at infinity, and for the line element 2.4

$$(2.12) \quad M = \lim_{r \rightarrow \infty} \frac{1}{8\pi} \int_{\mathcal{S}^2} \star d\xi = - \lim_{r \rightarrow \infty} \frac{1}{8\pi} \iint (r^2 \sin^2 \theta \tilde{\omega} \partial_r \tilde{\omega}) r^2 \sin \theta d\theta \wedge d\phi,$$

where $\xi = \xi_\mu dx^\mu$ and $\star$ is the Hodge operator. For r finite and sufficiently large the mass function is positive and behaves as $M(r) \sim 4a^4 / (9r^3) > 0$. The function tends to 0 as $r \rightarrow \infty$, which confirms our statement. The singular segment is the space-time singularity as curvature scalar is proportional to $\mathcal{D}$ which does not exist as a function on the segment, or more precisely, $\mathcal{D}$ is a distribution. Total angular momentum of an asymptotically flat space-time can be calculated as a surface integral over the sphere $\mathcal{S}^2$ at infinity, and for the line element 2.4

$$(2.13) \quad J = - \lim_{r \rightarrow \infty} \frac{1}{16\pi} \int_{\mathcal{S}^2} \star d\eta = - \lim_{r \rightarrow \infty} \frac{1}{16\pi} \iint (r^2 \sin^2 \theta \partial_r \tilde{\omega}) r^2 \sin \theta d\theta \wedge d\phi,$$

and we again obtain $J = a^2/3$ for solution 2.11. As we have mentioned earlier, the specific angular momentum for any regular element of the flow vanishes identically as trajectories of the flow are world-lines of locally non-rotating observers. Another way of seeing this is to calculate the form $d\star d\eta$, which measures angular momentum density when one take $dt = 0$, and which as a general form vanishes identically for $\tilde{\omega}$ satisfying equation 2.6. In the singular regions where $\tilde{\omega}$ is a distribution and does not satisfy 2.6, the contribution from $d\star d\eta$ to total angular momentum can be calculated from the above surface integral.

2.4. Asymptotically flat space-times of $\mathcal{K}$ flow are massless, some of them may have non-vanishing total angular momentum. The effect of vanishing of total mass should hold for all asymptotically flat space-times of $\mathcal{K}$ flow. Asymptotical flatness for line element 2.4 means that $\tilde{\omega}$ falls off like r^{-3} or quicker in spherical coordinates. If $\tilde{\omega} \sim r^{-3}$ then such space-time may have nonzero total angular momentum. Vanishing of total mass is due to the presence of singularities that contribute positive mass that cancel contribution from regular regions of the space-times where energy density is negative definite. Similarly, the existence of asymptotically flat $\mathcal{K}$ flows implies that the singularities they contain must have finite angular momentum equal to the total angular momentum of the space-times as the flow contributes no angular momentum in the regular regions.

3. CONCLUSIONS

We examined cylindrically symmetric and stationary dust flow along world-lines of locally non-rotating observers of which space trajectories are rings about common axis of rotation. Such flow can be constructed globally with the exception of regions where curvature singularities reside and that are additional sources of gravitational field. The flow is differential (the shear tensor is non-zero), non-expanding (the dilation scalar vanishes identically) and locally does not rotate (the vorticity tensor vanishes identically). Local mass density of the flow is necessarily negative-definite which makes the flow nonphysical as far as usual forms of matter are concerned.

There exist asymptotically flat space-times of the flow. Components of the flow move differentially on circular orbits with respect to an asymptotic stationary observer. The space-times have vanishing total mass and contain internal singularities

where distributional sources of positive mass and of angular momentum are located, and this should be understood in the following sense. It turns out that a surface integral over the sphere at infinity and which reproduces total mass (of the Kerr black hole, for example), vanishes for the solutions. For radii sufficiently large the resulting mass function is positive and attains zero monotonically in the limit of infinite radius. As local energy density is negative definite in the regions where space-time is smooth, the internal singularities of curvature the solutions contain, must be distributional sources of positive mass (maybe infinite) of which contribution to the total mass is screened by the regular regions, such the mass function is zero at infinity. This phenomenon is quite analogous with the screening of singularities of negative mass by regions of positive energy density of asymptotically flat van Stockum-Bonnor flow. By construction of $\mathcal{K}$ flow the specific angular momentum per particle is zero, nevertheless, total angular momentum of a class of the asymptotically flat space-times may be non-zero and we gave an example in 2.11. As the $\mathcal{K}$ flow has no specific angular momentum, the total angular momentum, which is given by a surface integral over the sphere at infinity and which reproduces total angular momentum (of the Kerr black hole, for example), must be located in singularities of the space-times.

Among other solutions, the model contains an infinite sequence of smooth asymptotically flat multipolar solutions to which a class of other asymptotically flat solutions can be decomposed. There also exist an infinite sequence of the corresponding internal multipolar solutions that are not asymptotically flat. The solutions are given in general by formulas 2.9 and 2.10.

4. ACKNOWLEDGEMENTS

This research is partially supported by the Polish Ministry of Science and Informatization, grants no. 1 P03D 005 28 and no. 1 P03B 012 29.

REFERENCES

- [1] W.J. van Stockum, *Proc. Roy. Soc. Edin.* **57**, 135 (1937)
- [2] J. M. Bardeen, *APJ*, **162**, 71 (1970)
- [3] W. B. Bonnor, *J. Phys. A: Math. Gen.* **10**, 10, 1673 (1977)
- [4] L. Bratek, J. Jalocha, M. Kutschera, *astro-ph/0603791*